

A 2 PN/RM metric of General Relativity

Olivier Minazzoli* and Bertrand Chauvineau
UNS, OCA-ARTEMIS UMR 6162

We present a derivation of 2PN/RM metric field equations from the Einstein field equation in General Relativity. We use the exponential parametrization and the c^{-3} isotropic spatial coordinates such as in IAU2000 recommendations.

Keywords: post newtonian relativistic motion; post newtonian; harmonic gauge; TIPO; ASTROD; laser telemetry in the solar system

I. INTRODUCTION

In a foreseeable future, telemetry will reach a new accuracy using laser links and high accuracy clocks (such as the TIPO experiments which is in study [1]). At such experiments accuracies, the effective 1,5 PN/RM (acronym defined in section II) metric defined from the 1PN IAU2000 [2] metric will be no longer enough to fit data (2 PN/RM effects lead to terms of about 10^{-10} s for a flying time of about 1000s when tagging units accuracy reach at least 10^{-11} s). Hence, it is necessary to develop the metric up to the 2 PN/RM order. Our work is completely based on General Relativity (GR) and hence we don't consider any alternative theory in this paper. This means that our result would have a practical usefulness in telemetry only if the GR theory is the "true" theory of gravity or if alternative theory deviates from GR only by sufficiently small post-Newtonian numerical parameters in the metric. This is compatible with present available data.

We start our work from the "exponential parametrization" such as in DSX [3] or in the IAU2000 resolutions [2]. Hence, our notations come from [3]. Other works on the subject have already been done ([4],[5]) but they didn't write their metric in the convenient exponential parametrization and these works have been done in harmonic gauge from the start to the end. This is not our case since we fix the gauge only at the end when we want peculiar solutions in some peculiar gauges.

In section II we introduce a new terminology before recalling some well known results from the DSX paper [3] in section III. Then we derive Einstein's equations in our metric's parametrization and finally we give formal solutions in any gauge that respect the exponential parametrization – with a particular attention to the harmonic gauge.

II. TERMINOLOGY : DEFINITION OF THE PN/BM AND PN/RM METRICS

The PN approximation is based on the assumption of a weak gravitational field and weak velocities (ie. of the

order $\sqrt{GM}/r$ or less, M being some characteristic mass of the system) for both the sources and the (test) body. It formally consists in looking for solutions under the form of an expansion in powers of $1/c$. The usually so-called n PN order terms in the metric, leading to c^{-2n} terms in the equation of motion of a body describing a bounded orbit, are terms of orders c^{-2n-2} in g_{00} , c^{-2n-1} in g_{0i} and c^{-2n} in g_{ij} . In this paper, a metric developed this way will be referred as the n PN/BM metric (BM meaning "Bounded Motion" for test particles). It is particularly well-adapted for studying bounded motions in systems made by non-relativistic massive bodies, as the Solar System is.

However, since we are interested in the propagation of light, we are lead to relax the hypothesis on the velocity of the test particle whose motion is considered. Of course, this doesn't change the full metric, but the terms to be considered in the metric components are not the same as in the PN/BM problem. Indeed, the terms leading to c^{-2n} terms in the equation of motion of a test particle moving with relativistic velocity are terms of order c^{-2n} in $g_{\alpha\beta}$ (ie. in both g_{00} , g_{0i} and g_{ij}). In this paper, a metric developed this way will be referred as the n PN/RM metric (RM meaning "Relativistic Motion" for test particles). A PN/RM metric is particularly well-suited for studying *relativistic* motions of test bodies (for instance, the propagation of light) in systems made by non-relativistic bodies, as the Solar System is.

PN/BM and PN/RM orders are illustrated in figure 1.

The present paper deals with the PN/RM problem since we are concerned in propagation of light.

III. 1,5 PN/BM METRIC

We recall here results from DSX[3] on the 1,5PN/BM metric. First, the "exponential parametrization" with the spatial isotropy condition makes the metric writes

$$g_{00} = -e^{-2w/c^2} \quad (1)$$

$$g_{0i} = -\frac{4}{c^3}w^i \quad (2)$$

$$g_{ij} = \gamma_{ij}e^{2w/c^2}. \quad (3)$$

*also in the IRAP PhD program

IV. 2 PN/RM METRIC




FIG. 1: General scheme of orders taken into account in PN/BM and PN/RM metrics.

With $\gamma_{ij} = \delta_{ij} + O(c^{-4})$. From which follows

$$g^{00} = -e^{2w/c^2} + O(c^{-6}) \quad (4)$$

$$g^{0i} = -\frac{4}{c^3}w_i + O(c^{-5}) \quad (5)$$

$$g^{ij} = \delta^{ij}e^{-2w/c^2} + O(c^{-4}). \quad (6)$$

The Einstein field equation then reduces to four equations on w and w_i

$$\Delta w + \frac{3}{c^2}\partial_t^2 w + \frac{4}{c^2}\partial_t\partial_j w_j = -4\pi G\sigma + O(c^{-4}) \quad (7)$$

$$\Delta w_i - \partial_{ij}^2 w_j - \partial_t\partial_i w = -4\pi G\sigma^i + O(c^{-2}). \quad (8)$$

Where $\sigma = c^{-2}(T^{00} + T^{kk})$, $\sigma^i = c^{-1}T^{0i}$ and $\Delta = \vec{\partial} \cdot \vec{\partial}$. Hence, to the 1,5PN/BM approximation correspond 4 equations of the GR equation and the gauge invariance keep 1 degree of freedom (indeed σ^μ is approximately divergence free). This gauge invariance is characterized by the arbitrary differentiable function λ in DSX.

$$w' = w - \frac{1}{c^2}\partial_t\lambda \quad (9)$$

$$w'_i = w_i + \frac{1}{4}\partial_i\lambda. \quad (10)$$

The IAU2000 resolution then corresponds to a peculiar choice of gauge in order to fix the last degree of freedom : the harmonic gauge. This condition writes, in the 1,5PN/BM approximation $\{g^{\alpha\beta}\Gamma_{\alpha\beta}^0 = O(c^{-5}), g^{\alpha\beta}\Gamma_{\alpha\beta}^i = O(c^{-4})\}$. The four metric field equations then reduce to

$$\square_m w = -4\pi G\sigma + O(c^{-4}) \quad (11)$$

$$\Delta w_i = -4\pi G\sigma^i + O(c^{-2}) \quad (12)$$

where $\square_m = \Delta - \partial_t^2$.

At the 2PN/RM order, we get 6 new degrees of freedom in the metric. As we know, from the 10 degrees of freedom in the equation of GR, there are only 6 cleared by the 10 equations since 4 degrees of freedom correspond to a gauge invariance. Indeed, the metric is determined by the field equations modulo the knowing of the used coordinates. Since, there are four coordinates, there are 4 degrees of freedom not reduced by the 10 field equations. As seen in the previous section, to the 1,5PN/BM approximation correspond 4 metric field equations and the gauge invariance, characterized by λ , keep 1 degrees of freedom. Then, we expect from the 2PN/RM order 6 equations with a gauge invariant field (characterized by a 3-vector) which (the equations) reduce the 6 degrees of freedom to 3. Let us fix the covariant form of the metric

$$g_{00} = -e^{-2w/c^2} \quad (13)$$

$$g_{0i} = -\frac{4}{c^3}w^i \quad (14)$$

$$g_{ij} = \delta_{ij}e^{2w/c^2} + \frac{4\tau_{ij}}{c^4} + O(c^{-5}) \\ = \delta_{ij} \left(1 + \frac{2w}{c^2} + \frac{2w^2}{c^4} \right) + \frac{4\tau_{ij}}{c^4} + O(c^{-5}). \quad (15)$$

Which implie

$$g^{00} = -e^{2w/c^2} + O(c^{-6}) \quad (16)$$

$$g^{0i} = -\frac{4}{c^3}w_i + O(c^{-5}) \quad (17)$$

$$g^{ij} = \delta^{ij}e^{-2w/c^2} - \frac{4\tau_{ij}}{c^4} + O(c^{-5}). \quad (18)$$

Then we can derive the 2PN/RM field equations from the Einstein field equation $R^{\alpha\beta} = \frac{8\pi G}{c^4}(T^{\alpha\beta} - \frac{1}{2}g^{\alpha\beta}g_{\rho\sigma}T^{\rho\sigma})$.

$$\frac{2}{c^4}\Theta_{ij}(\tau_{kl}) + \Xi_{ij}(w, w_k) = \\ \frac{8\pi G}{c^4} \left(\frac{1}{2}\eta^{ij}T^{00} + T^{ij} - \frac{1}{2}\eta^{ij}T^{kk} - \frac{2w}{c^2}\eta^{ij}T^{00} \right). \quad (19)$$

Where

$$\Theta_{ij}(\tau_{kl}) = \partial_{ki}^2\tau_{kj} + \partial_{kj}^2\tau_{ik} - \Delta\tau_{ij} - \partial_{ij}^2\tau_{kk}, \quad (20)$$

$$\Xi_{ij}(w, w_k) = \\ -\frac{\delta_{ij}}{c^2}\square_m w + \frac{2}{c^3}(\partial_{0i}^2 w_j + \partial_{0j}^2 w_i) - \frac{2}{c^4}\partial_i w \partial_j w + \frac{4\delta_{ij}}{c^4}w\Delta w. \quad (21)$$

Then, considering w and w_k as source terms and identifying the relation $-\frac{c^4}{2}\left(\frac{4\delta_{ij}}{c^4}w\Delta w\right) = \frac{8\pi G}{c^2}\eta^{ij}wT^{00} + O(c^{-2})$ from (11), we get

$$\Theta_{ij}(\tau_{kl}) = 4\pi G\sigma^{ij} - [(\partial_{ti}^2 w_j + \partial_{tj}^2 w_i - \partial_i w \partial_j w) + 2\eta^{ij}(\partial_t^2 w + \partial_{tk}^2 w_k)] \quad (22)$$

Where $\sigma^{ij} = T^{ij} - \eta^{ij}T^{kk}$. As expected, the operator Θ is gauge invariant through a relation on any 3-vector field. Indeed, we have $\Theta_{ij}(\tau_{kl} + \partial_k A_l + \partial_l A_k) = \Theta_{ij}(\tau_{kl})$, whatever is the 3-vector field A_k . Hence, the 2PN/RM gauge invariance is characterized by both the arbitrary differentiable function λ and the arbitrary differentiable 3-vector A_k .

$$w' = w - \frac{1}{c^2} \partial_t \lambda \quad (23)$$

$$w'_i = w_i + \frac{1}{4} \partial_i \lambda \quad (24)$$

$$\tau'_{kl} = \tau_{kl} + \partial_k A_l + \partial_l A_k + \frac{1}{2} \delta_{kl} \partial_t \lambda. \quad (25)$$

A. General formal solution

We can show that it always exists a gauge transformation $\tau'_{kl} = \tau_{kl} + \partial_k A_l + \partial_l A_k$ that reduces the operator Θ to a Laplacian ($\Theta_{ij}(\tau'_{kl}) = -\Delta \tau'_{ij}$). To be placed on this gauge, A_k must satisfy $\Delta A_j = \partial_k \tau_{kj} - \frac{1}{2} \partial_j \tau_{kk}$ which is clearly inversible since we know the laplacian's green function. Hence, there always exists a peculiar gauge where the solution is

$$\tau_{ij} = \Delta^{-1} \{ -4\pi G\sigma^{ij} + [(\partial_{ti}^2 w_j + \partial_{tj}^2 w_i - \partial_i w \partial_j w) + 2\eta^{ij}(\partial_t^2 w + \partial_{tk}^2 w_k)] \}, \quad (26)$$

where

$$\Delta^{-1} \{ f(t, \vec{x}) \} = \int \frac{f(t, \vec{x}')}{|\vec{x} - \vec{x}'|} d^3 x'. \quad (27)$$

Then, from this equation, we can get the solution τ_{kl} in any gauge, using any vector field V_k with $\tau'_{kl} = \tau_{kl} + \partial_k V_l + \partial_l V_k$.

B. The harmonic gauge

The harmonic gauge condition stands $g^{\alpha\beta}\Gamma_{\alpha\beta}^\gamma = 0$ (Or $\partial_\sigma(\sqrt{-g}g^{\sigma\gamma}) = 0$ with g the metric determinant). As seen in DSX [3], the harmonic 1,5PN/BM condition $g^{\alpha\beta}\Gamma_{\alpha\beta}^0 = O(c^{-4})$ gives the usual harmonic constraint on the "potentials" ($\partial_t w + \partial_k w_k = O(c^{-2})$). While, the conditions $g^{\alpha\beta}\Gamma_{\alpha\beta}^i = O(c^{-4})$ give no more information due to the "isotropy condition" which makes spatial coordinates always harmonic modulo $O(c^{-4})$. At the 2PN/RM

level, the harmonic constraint is $g^{\alpha\beta}\Gamma_{\alpha\beta}^\gamma = O(c^{-5})$. That gives the harmonic conditions on the "potentials" at the 2PN/RM level

$$\partial_t w + \partial_k w_k = O(c^{-2}) \quad (28)$$

$$\partial_k \tau_{ik} - \frac{1}{2} \partial_i \tau_{kk} + \partial_t w_i = O(c^{-1}). \quad (29)$$

Then $\Theta_{ij}(\tau_{kl})$ writes:

$$\Theta_{ij}(\tau_{kl}) = -\{\Delta \tau_{ij} + \partial_t(\partial_i w_j + \partial_j w_i)\}. \quad (30)$$

Hence, considering w and w_i as source terms, the solution of the Einstein field equations in the harmonic gauge is :

$$\tau_{ij} = \Delta^{-1} \{ -4\pi G\sigma^{ij} - \partial_i w \partial_j w \}, \quad (31)$$

where w and w_j satisfy the 1,5PN/BM harmonic condition (28).

from non-harmonic to harmonic gauge

If τ_{ij} is not an harmonic solution, then the gauge transformation $\tau_{ij} = \epsilon_{ij} + \partial_i A_j + \partial_j A_i$ makes ϵ_{ij} an harmonic solution if A_j satisfies

$$\Delta A_j = \partial_k \tau_{kj} - \frac{1}{2} \partial_j \tau_{kk} + \partial_t w_j \quad (32)$$

where w_j must be expressed in the harmonic gauge with the 1,5PN/BM field equations (12). Since it is inversible, this relation always allow us to put the metric in the harmonic gauge, whatever is the gauge from which we start.

V. CONCLUSIONS

In order to fit future inter-planetary telemetry data, we developed a GR metric up to the 2 PN/RM order with the DSX "exponential parametrization". We gave both general and harmonic formal solutions. A more detailed version of this result can be found in [6]. An explicit solution of these equations and of the (non-massive-)Scalar-Tensor 2PN/RM field equations in the case of the solar system is in preparation.

Acknowledgments

Olivier Minazzoli wants to thank the Government of the Principality of Monaco for their financial support.

-
- [1] Samain, E., Bonnefond, P., Nicolas, J. 2003 Phys. & Chem. of the Earth
[2] Soffel, M. & al. 2003AJ....126.2687S
[3] Damour, T., Soffel, M., Xu, C. 1991PhRvD..43.3273D
[4] Blanchet, L. 1996PhRvD..54.1417B
[5] Anderson, J.L., Decanio, T.C. 1975GReGr...6..197A
[6] Minazzoli O., Chauvineau, B. 2009PhRvD..79h4027M